

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> :

C12N 15/31, 15/44, 15/62

C07K 13/00, A61K 39/02

C12N 1/20 // C12P 21/08

A1

(11) International Publication Number:

WO 93/04175

(43) International Publication Date:

4 March 1993 (04.03.93)

(21) International Application Number: PCT/EP92/01827

(22) International Filing Date: 11 August 1992 (11.08.92)

(30) Priority data:

9117602.4	15 August 1991 (15.08.91)	GB
9122301.6	21 October 1991 (21.10.91)	GB
9211318.2	28 May 1992 (28.05.92)	GB
9211317.4	28 May 1992 (28.05.92)	GB

(71) Applicants (for all designated States except US): SMITH-KLINE BEECHAM BIOLOGICALS (S.A.) [BE/BE]; 89, rue de l'Institut, B-1330 Rixensart (BE). MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Bunsenstrasse 10, D-3400 Göttingen (DE). DEUTSCHES KREBSFORSCHUNGSZENTRUM STIFTUNG DES ÖFFENTLICHEN RECHTS [DE/DE]; Im Neuenheimer Feld 280, D-6900 Heidelberg 1 (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LOBET, Yves [BE/BE]; SmithKline Beecham Biologicals (S.A.), 89, rue de l'Institut, B-1330 Rixensart (BE). SIMON, Markus [DE/DE]; SCHAIBLE, Ulrich [DE/DE]; Max-Planck-Institut für Immunobiologie, D-7800 Freiburg (DE). WALLICH, Reinhard [DE/DE]; Deutsches Krebsforschungszentrum Stiftung des öffentlichen Rechts, Unter den Eichen 6, D-6900 Heidelberg (DE). KRAMER, Michael [DE/DE]; Max-Planck-Institut für Immunobiologie, D-7800 Freiburg (DE).

(74) Agent: DALTON, Marcus, Jonathan, William; Smith-Kline Beecham, Corporate Patents, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB).

(81) Designated States: AU, CA, JP, KR, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).


Published

With international search report.

(54) Title: OSP A PROTEINS OF BORRELIA BURGDORFERI SUBGROUPS, ENCODING GENES AND VACCINES

## (57) Abstract

The invention relates to novel Borrelia, and OspA antigens derived therefrom. These antigens show little homology with known OspA's and are therefore useful as vaccine and diagnostic reagents. Multicomponent vaccines based on OspA's from different Borrelia groups are also disclosed.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain			US	United States of America

-1-

## OSP A PROTEINS OF BORRELIA BURGDORFERI SUBGROUPS, ENCODING GENES AND VACCINES.

The present invention relates to novel antigens, to methods for their production, to compositions containing them and to their use in the prevention, treatment and diagnosis of Lyme Disease in humans and other animals. In particular the present invention discloses novel serotypes/genotypes of the outer surface protein (Osp A) from the spirochete Borrelia burgdorferi, the causative agent of Lyme disease, and vaccine and diagnostic reagents based on B. burgdorferi of more than one subgroup.

Lyme disease in humans is a chronic progressive disease caused by B. burgdorferi, which is transmitted to humans mainly by Ixodes ticks. The disease attacks many organs, notably the skin, heart, liver, central and peripheral nervous system, kidneys as well as the musculoskeletal system.

Lyme disease itself is the most common vector borne infection in the USA and has been reported in every continent except Antarctica.

A number of groups have isolated and proposed the major surface protein (Osp A) of B. burgdorferi, as being a potential vaccine candidate for use against Lyme disease. For example, International patent application published under WO90/04411 (SIMBICOM) discloses the cloning and expression of an Osp A protein derived from B. burgdorferi B31 and its use as a vaccine, M.M. Simon and colleagues have cloned and expressed Osp A from B. burgdorferi ZS7 (European Patent Application No. 90117943.2 published under No. 0418 827), and demonstrated its protective capacity to induce antibodies in passively immunised SCID mice (PNAS : 87 1990, 3768 - 3772). Flavell and colleagues (Science (1991) 250 p553-556) have cloned and expressed the gene for Osp A from B. burgdorferi, N40 and have demonstrated its protective efficacy in C3H/He mice.

All isolates of B. burgdorferi identified above appear to be closely related. However we have now identified six subgroups of B. burgdorferi by analysing 55 spirochete isolates from different geographical areas and sources with series of immunological, biochemical and molecular genetic techniques. The finding of different subgroups of B. burgdorferi isolates

-2-

have important implications for both effective vaccination and diagnosis of Lyme disease. Since ELISA diagnostic assays directed against a strain of which species N40 is a member would not or only partially cross react with species from other subgroups. Equally a vaccine based on only one Osp A for example from N40 would not provide optimal protection against B. burgdorferi from a different subgroup.

The present inventors have identified five additional subgroups of B. burgdorferi based on their Southern blotting and PCR amplification of Osp A sequences, and differences in the amino acid and nucleic acid sequences themselves and also in reactivity against monoclonal antibodies to Osp A proteins.

The surprising discovery that B. burgdorferi exhibits such heterogeneity has important implications for vaccines against, and diagnostics reagents for the detection of, Lyme disease, since vaccines or diagnostic based on an Osp A from one group of B. burgdorferi may not detect or protect against infection of B. burgdorferi from a second group. Indeed the present inventors have shown that whereas protection can be afforded by anti Osp A antibodies generated by recombinant Osp A or viable or killed organisms within a group of closely related strains, no or only partial, protection is observed if challenge is made with an organism from a different group.

A first subtype having representative ZS7, B31, N40 is referred to herein as group I (alternative nomenclature refers to group I strains as group A).

A second subgroup hereinafter group II (alternatively known as group B) is herein disclosed and is exemplified by the species ZQ1. This species has been deposited at the DSM Deutsche Sammlung von Mikroorganismen Und Zellkulturen of Mascheroderweg 91B D-3300 Braunschweig on 11 July 1991 and given the accession No. DSM6606. This group is distinct in a number of ways from group I.

Firstly, plasmid analysis of strain ZQ1 when compared with representatives from group I shows that ZQ1 has at least two unique plasmids of 18 and 14Kd. Furthermore plasmid of 16Kb found in group I strains is absent from strain ZQ1. Secondly when examined by

-3-

polyacrylamide gel electrophoresis Osp A from ZQ1 has an apparent molecular weight of 32Kd.

Thirdly various monoclonal antibodies specific for Osp A of group I, such as the antibodies designated LA2 LA26 LA28 LA33 do not react with Osp A from group II. LA2 and LA26 antibody are known and have been described in European patent application No. EP0418827. The hybridoma producing LA2 has been deposited under accession No. ECACC 89 09 1302 on 13th September 1989 at the European Collection of animal cell cultures, Public health laboratory services Porton Down Wiltshire SP40J9. The hybridoma producing LA26 has been deposited at the same culture collection under accession No. 9005406 on 28th June 1990.

However an other monoclonal antibody, LA31.1, does react with the Osp A species of subgroups I and II suggesting common epitopes among Osp A species of different subgroups do exist.

We have also characterised Osp A DNA from group II strains by PCR amplification. Two pairs of Osp A primers were used, these being designated prOsp A1-prOsp A4 and prOsp A1-prOsp A2.

The primers have the following sequences:

prOsp A1 : 5'-GGGAATAGGT CTAATATTAG CC-3'  
prOsp A2 : 5'-TGCCTGAATT CCAAGCTGCA-3'  
prOsp A4 : 5'-GCAGTTAAAG TTCCTTCAAG AACATAGC-3'

and correspond to nucleotides at positions 138-160 (prOsp A1), 611-638 (prOsp A4), and 759-778 (prOsp A2), of ZS7 Osp A respectively. Total DNA of Group II, B. burgdorferi are distinguishable from that of Group I in that group II will not permit successful PCR (polymerase chain reaction) amplification when prOsp A1 and prOsp A2 are used as primers whereas total DNA from group I will permit a successful reaction. In contrast, when prOsp A1 and prOsp A4 are used as primers, then both DNA's of group I and group II will undergo polymerase chain reaction.

These results show that group II, B. burgdorferi have a different Osp A sequences to group I Osp As.

Equally, group II Osp A DNA, can be distinguished from group I Osp A DNA by Southern blotting utilising Osp A ZS7 sequence as a probe. Digesting genomic DNA of B. burgdorferi group II with the restriction enzyme Hind III shows a different banding pattern than that of group I strain.

For example a group I strain reveals two hybridisation fragments of 1.2 kb and 0.3 kb whereas a group II strain expressed two fragments of 0.9 kb and 0.4 kb.

Also, passive immunisation of SCID mice with immune sera to Osp A from group II will not protect against B. burgdorferi group I challenge.

Finally when compared to the sequence of Osp A ZS7 the DNA and amino acid sequence of ZQ1 strain Osp A Sequence ID 1 and 2 (i.e. a group II strain), demonstrates the substantial variation between a group I and group II strain at both the DNA and amino acid levels.

The surprising inability to protect against heterologous challenge means that a vaccines based on a single Osp A from a group I organism may be ineffective against infection with B. burgdorferi from group II organism and vice versa. Consequently there is a need for a vaccine which will protect against group II infection.

Accordingly there is provided DNA sequence encoding an Osp A derived from a B. burgdorferi characterised in that polymerase chain reaction amplification will take place utilising prOsp A1-A4 primers, but not prOsp A1-A2, and the invention further provides purified or isolated Osp A encoded thereby.

Preferably the Osp A of the present invention will not react with monoclonal antibody LA2 or LA26. Most preferably the Osp A of the present invention will have a relative molecular weight of 32 kDa as determined by SDS electrophoresis utilising molecular weight markers as standard.

Preferably th Osp A is derived from either ZQ1 or NE11H or other group

-5-

II strain.


Preferably the protein is at least 70% pure as determined by SDS polyacrylamide gel electrophoresis, in particular at least 80% pure, and most preferably at least 95% pure. Preferably the Osp A antigen has a relative molecular weight of 32kDa as determined by SDS polyacrylamide gel electrophoresis. Preferably the Osp A antigen will not react with the monoclonal antibody LA2.

10 In another aspect of the present invention there is provided a DNA sequence substantially identical to the DNA sequence encoding Osp A from ZQ1. By substantially identical it is meant a DNA sequence contains a sequence which is at least 85% identical and preferably 90% identical to the ZQ1 Osp A sequence depicted in Sequence ID 1. The mature protein starts at position 17 and preferably the invention provides a DNA sequence which is at least 95% identical to the sequence encoding the mature protein as indicated in Sequence ID 2.

20 In an alternative embodiment there is provided an Osp A containing an amino acid sequence substantially identical to the sequence set forth in sequence ID 2. The term substantially identical in relation to the amino acid sequence means amino acid sequence which is at least 85% and preferably 90% identical to the protein sequence depicted in figure and preferably at least 85% identical to the mature protein.

25 The present inventors have also found utilising the same biochemical and immunological techniques other groups of *B. burgdorferi* unrelated to other group I or II. Two isolates characteristic of group III (or group C) are the strain designated as 19857 and 21038. Such strains, like group II, strains react under PCR conditions with prOsp A1-A2 but not with prOsp A1-A4. However they differ from group II strains when analysed by Southern blotting when utilising a ZS7 probe and restriction enzyme HindIII. Such analysis reveal hybridisation fragments of 4.0kb and 0.5kb. In addition they react with monoclonal antibody LA26, but not LA2.

35 We have now sequenced utilising PCR technology, and expressed a DNA sequence encoding an Osp A from strain 19857. The DNA and amino acid sequence are depicted in sequence ID No. 3 and 4. A comparison with





-6-

sequence from other groups shows that the Osp A only has approximately 65% identity at the DNA and amino acid levels with Osp A from other subgroups.

- 5 Accordingly the present invention provides a DNA sequence encoding an Osp A protein wherein the DNA sequence is substantially identical to the DNA sequence of Sequence ID 3. In a further embodiment there is provided an Osp A protein which has an amino acid sequence substantially identical to the protein as disclosed in Sequence ID 4.

10

The term substantial identical as used herein with respect to the sequences depicted in sequence ID No.3 and 4 mean at least 70% identity, preferably 80 - 85% identity, more preferably 85 - 95% identical.

- 15 A fourth subgroup, group IV (or group D) of B. burgdorferi, as exemplified by a Swedish isolate ACA-1 differs from any of groups I, II or III, M. Jonsson et al. (Int. Conf. Lyme Borreliosis 1990). This group will not undergo PCR with either prOsp A1-A2 or prOsp A1-A4. Their Osp A's react with monoclonal antibody LA26 but not with LA2.

20

Southern blotting analysis when utilising a ZS7 probe and restriction enzyme HindIII reveal a single hybridisation fragment of about 1.7 kb.

- Isolated or recombinant Osp A from group IV also form part of the present invention as do vaccine or diagnostic compositions containing them.

25

- By utilising the same Biochemical techniques, the present inventors have identified representatives from group V and group VI. A group V isolate is known as 20047 and a representative of group VI is the strain known as

30

S90.

Isolated Osp A from a group V and VI strain also form part of the invention.

- 35 The present invention also relates to immunogenic derivatives of the OspA from group II, group III, group IV and group V or group VI B. burgdorferi.

The term immunogenic derivative as used herein encompasses any

-7-

mol cule such as a truncated or hybrid protein which are immunologically reactive with antibodies generated by infection of a mammalian host with a group II, or group III or group IV, or group V or group VI, B. burgdorferi and which preferably do not react with the monoclonal antibodies designated LA2. Such derivatives may be prepared by substitution, or rearrangement of aminoacids or by chemical modifications thereof.

Immunogenic fragments of the protein, which may be useful in the preparation of subunit vaccines, may be prepared by expression of the appropriate gene fragments or by peptide synthesis, for example using the Merrifield synthesis (The Peptides, Vol 2., Academic Press, NY, page 3).

The immunogenic derivative of the invention can be a hybrid, that is, a fusion polypeptide containing additional sequences which can carry one or more epitopes for other B. burgdorferi immunogens including Osp A and/or Osp B, or other immunogens from, for example other pathogens. Alternatively, the immunogenic derivative of the invention can be fused to a carrier polypeptide such Hepatitis B surface or core antigen or to another carrier which has immunostimulating properties, as in the case of an adjuvant, or which otherwise enhances the immune response to the Osp A protein or derivative thereof, or which is useful in expressing, purifying or formulating the protein or derivative thereof.

The invention also extends to the Osp A protein or immunogenic derivative thereof when chemically conjugated to a macromolecule using a conventional linking agent such as glutaraldehyde (Geerlings et al, (1988) J. Immunol. Methods, 106, 239-244).

Particularly preferred derivatives are hybrid fusion proteins of the influenza non-structural protein NS1 and Osp A or truncated Osp A derivatives. These can advantageously be produced by recombinant DNA techniques in high yields in E. coli.

Accordingly the present invention provides an Osp A protein fused to the N terminal portion of the protein NS1 from Influenza. Preferably the Osp A portion of the fusion protein is truncated so as to remove at least the hydrophobic signal sequence. More preferably the N terminal 16 amino acids representing the signal sequence and the cysteine residue which is

-8-

the first amino-acid of the mature protein of the Osp A are missing.

Preferably the NS1 part of the fusion protein comprises from 3 to 81 amino acids of the N-terminal portion of the gene. The longer the NS1 part of the fusion is preferred, since this permits higher expression levels of proteins.

These hybrid proteins of the present invention are recoverable from the soluble fraction as non aggregated protein, in contrast to the native recombinant Osp A and are expressed at a higher level. The proteins retain their ability to react with monoclonal antibodies raised against native Osp A and are capable of raising a protective immune response in laboratory animals eg mice and rabbits.

In an embodiment of the invention there is provided an NS1 Osp A fusion protein from the Osp A group, known as group I, eg an Osp A from strain ZS7 as disclosed in EP 0418 - 827, or B31 as described by Howe *et al.* Science 227 p645, 1985, or N40 as described by Flavell *et al.* Such Osp A's show significant sequence homology at both the DNA and amino acid levels and are cross protective.

In a further embodiment of the present invention there is provided an NS1 - Osp A wherein the Osp A is from the group known as group II. For example the Osp from B. burgdorferi strain ZQ1 as described herein.

In a yet further embodiment of the present invention there is provided an NS1 - Osp A wherein the Osp A is from the group known as group III. For example strain 19857. The invention further provides a group IV NS1-Osp A, for example wherein the Osp A is derived from the strain AcaI.

The present invention further provides a DNA sequence which encodes an NS1- Osp A fusion protein as described herein. In particular the present invention provides a DNA sequence encoding an NS1 - Osp A protein and the protein itself as shown in Sequence ID Nos 5 to 12 attached herewith.

The proteins of the present invention can be produced by methods standard in the art for producing other Osp A proteins from known

-9-

strains. For example, the Osp A proteins may be produced in E. Coli utilising recombinant DNA technology. In particular, the proteins may be produced as full length immature proteins or mature proteins. The proteins may also be expressed as fusion proteins, such as NS1-fusion proteins.

A DNA sequence encoding the proteins of the present invention (eg Osp A ZQ1 or Osp A 19857) can be synthesized using standard DNA synthesis techniques, such as by enzymatic ligation as described by D.M. Roberts *et al* in Biochemistry 1985, 24, 5090-5098, by chemical synthesis, by *in vitro* enzymatic polymerization, or by PCR technology utilising for example a heat stable polymerase, or by a combination of these techniques.

Enzymatic polymerisation of DNA may be carried out *in vitro* using a DNA polymerase such as DNA polymerase I (Klenow fragment) in an appropriate buffer containing the nucleoside triphosphates dATP, dCTP, dGTP and dTTP as required at a temperature of 10<sup>0</sup>-37<sup>0</sup>C, generally in a volume of 50µl or less. Enzymatic ligation of DNA fragments may be carried out using a DNA ligase such as T4 DNA ligase in an appropriate buffer, such as 0.05M Tris (pH 7.4), 0.01M MgCl<sub>2</sub>, 0.01M dithiothreitol, 1mM spermidine, 1mM ATP and 0.1mg/ml bovine serum albumin, at a temperature of 4<sup>0</sup>C to ambient, generally in a volume of 50ml or less. The chemical synthesis of the DNA polymer or fragments may be carried out by conventional phosphotriester, phosphite or phosphoramidite chemistry, using solid phase techniques such as those described in 'Chemical and Enzymatic Synthesis of Gene Fragments - A Laboratory Manual' (ed. H.G. Gassen and A. Lang), Verlag Chemie, Weinheim (1982), or in other scientific publications, for example M.J. Gait, H.W.D. Matthes, M. Singh, B.S. Sproat, and R.C. Titmas, Nucleic Acids Research, 1982, 10, 6243; B.S. Sproat and W. Bannwarth, Tetrahedron Letters, 1983, 24, 5771; M.D. Matteucci and M.H. Caruthers, Tetrahedron Letters, 1980, 21, 719; M.D. Matteucci and M.H. Caruthers, Journal of the American Chemical Society, 1981, 103, 3185; S.P. Adams *et al.*, Journal of the American Chemical Society, 1983, 105, 661; N.D. Sinha, J. Biernat, J. McMannus, and H. Koester, Nucleic Acids Research, 1984, 12, 4539; and H.W.D. Matthes *et al.*, EMBO Journal, 1984, 3, 801.

Alternatively, the coding sequence can be derived from B. burgdorferi

-10-

mRNA, using known techniques (e.g. reverse transcription of mRNA to generate a complementary cDNA strand), and commercially available cDNA kits.

- 5 DNA polymers which encode mutants of the Osp A proteins may be prepared by site-directed mutagenesis of the cDNA which codes for the protein by conventional methods such as those described by G. Winter *et al* in Nature 1982, 299, 756-758 or by Zoller and Smith 1982; Nucl. Acids Res., 10, 6487-6500, or deletion mutagenesis such as described by Chan  
10 and Smith in Nucl. Acids Res., 1984, 12, 2407-2419 or by G. Winter *et al* in Biochem. Soc. Trans., 1984, 12, 224-225.

The process of the invention may be performed by conventional recombinant techniques such as described in Maniatis *et. al.*, Molecular  
15 Cloning - A Laboratory Manual; Cold Spring Harbor, 1982-1989.

In particular, the process may comprise the steps of:

- 20 i) preparing a replicable or integrating expression vector capable, in a host cell, of expressing a DNA polymer comprising a nucleotide sequence that encodes the protein or an immunogenic derivative thereof;
- 25 ii) transforming a host cell with said vector;
- iii) culturing said transformed host cell under conditions permitting expression of said DNA polymer to produce said protein; and
- 30 iv) recovering said protein.

The term 'transforming' is used herein to mean the introduction of foreign DNA into a host cell. This can be achieved for example by transformation, transfection or infection with an appropriate plasmid or viral vector using  
35 e.g. conventional techniques as described in Genetic Engineering; Eds. S.M. Kingsman and A.J. Kingsman; Blackwell Scientific Publications; Oxford, England, 1988. The term 'transformed' or 'transformant' will hereafter apply to the resulting host cell containing and expressing the

-11-

foreign gene of interest.

The expression vectors are novel and also form part of the invention.

- 5 The replicable expression vectors may be prepared in accordance with the invention, by cleaving a vector compatible with the host cell to provide a linear DNA segment having an intact replicon, and combining said linear segment with one or more DNA molecules which, together with said linear segment encode the desired product, such as the DNA polymer encoding  
10 the Osp A protein, or derivative thereof, such as an NS1 Osp A-fusion protein under ligating conditions.

Thus, the DNA polymer may be preformed or formed during the construction of the vector, as desired.

15

The choice of vector will be determined in part by the host cell, which may be prokaryotic or eukaryotic. Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses.

- 20 The preparation of the replicable expression vector may be carried out conventionally with appropriate enzymes for restriction, polymerisation and ligation of the DNA, by procedures described in, for example, Maniatis *et al.* cited above.

- 25 The recombinant host cell is prepared, in accordance with the invention, by transforming a host cell with a replicable expression vector of the invention under transforming conditions. Suitable transforming conditions are conventional and are described in, for example, Maniatis *et al.* cited above, or "DNA Cloning" Vol. II, D.M. Glover ed., IRL Press Ltd,  
30 1985.

- The choice of transforming conditions is determined by the host cell. Thus, a bacterial host such as *E. coli* may be treated with a solution of  $\text{CaCl}_2$  (Cohen *et al.*, Proc. Nat. Acad. Sci., 1973, 69, 2110) or with a  
35 solution comprising a mixture of  $\text{RbCl}$ ,  $\text{MnCl}_2$ , potassium acetate and glycerol, and then with 3-[N-morpholino]-propane-sulphonic acid,  $\text{RbCl}$  and glycerol. Mammalian cells in culture may be transformed by calcium co-precipitation of the vector DNA onto the cells. The invention also

-12-

extends to a host cell transformed with a replicable expression vector of the invention.

5 Culturing the transformed host cell under conditions permitting expression of the DNA polymer is carried out conventionally, as described in, for example, Maniatis *et al* and "DNA Cloning" cited above. Thus, preferably the cell is supplied with nutrient and cultured at a temperature below 50°C.

10 The product is recovered by conventional methods according to the host cell. Thus, where the host cell is bacterial, such as *E. coli* it may be lysed physically, chemically or enzymatically and the protein product isolated from the resulting lysate. Where the host cell is mammalian, the product  
15 extracts. Conventional protein isolation techniques include selective precipitation, adsorption chromatography, and affinity chromatography including a monoclonal antibody affinity column.

20 The proteins of the present invention when expressed in *E. Coli* as full length proteins produce lipoprotein micelles which are immunogenic. Such full length lipoproteins form part of the present invention.

It will be appreciated that all the proteins of the present invention may find utility in vaccine and diagnostic applications.

25 Furthermore, it will be appreciated that other antigenic components such as other Osp A antigens or Osp B from *B. burgdorferi* isolates or antigens from other pathogens may be included in a vaccine composition of the present invention. Additionally other antigens from other organisms may  
30 be present in the vaccine.

The Osp A antigens utilised in the compositions of the present invention include immunogenic fragments of Osp A, i.e. fragments of Osp A containing immunogenic B or T-cell epitopes, or fusion proteins containing  
35 such epitopes, including fusions of Osp A's from different groups.

In an aspect of the present invention there is provided a vaccine or diagnostic composition comprising an Osp A or derivative thereof from a

-13-

- subgroup II strain. In particular on characteris d in that it is encoded by a DNA sequence which will permit successful polymerase chain reaction to take place utilising prOsp A1-A4 primers, but not prOsp A1-A2, the protein being in admixture with suitable excipient or carrier. More
- 5 particularly there is provided a composition comprising an Osp A or derivative thereof from a group II strain having an amino acid sequence substantially identical to the protein sequence depicted in sequence ID No.2.
- 10 Such compositions may advantageously also contain an Osp A derivative from a subgroup I strain.

Accordingly the present provides in a preferred embodiment a composition, preferably a vaccine composition comprising an Osp A

15 antigen derived from a subgroup I B. burgdorferi, and an Osp A antigen from a subgroup II, B. burgdorferi.

Preferably the Osp A antigen from group I are selected from group, B31, ZS7 and N40.

20 Preferably the Osp A antigen from group II is selected from the group ZQ1, NE11H.

The present invention also provides a vaccine or diagnostic composition

25 comprising an OspA or derivative from a subgroup III strain, the protein being a admixture with a suitable excipient or carrier. More particularly the vaccine or diagnostic composition comprises an OspA or derivative thereof having an amino acid sequence substantially identical to the protein sequence depicted in sequence ID No.4.

30 Preferably the OspA is derived from strain 19857.

The invention also provides a vaccine or diagnostic composition comprising an Osp A or derivative from subgroup IV, V or VI.

35 In a preferred embodiment there is provided a combination vaccine comprising at least two OspA's or derivatives thereof from different B. burgdorferi subgroups. Such vaccine compositions provide protection from



-14-

a wider range of *Borrelia* infections than hitherto. Preferably the vaccine comprises an Osp A from each of groups I to IV and most preferably an Osp A from each of group I to VI.

5 In the vaccine of the invention, an aqueous solution of the protein(s) can be used directly. Alternatively, the protein, with or without prior lyophilization, can be mixed or absorbed with any of the various known adjuvants. Such adjuvants include, but are not limited to, aluminium hydroxide, muramyl dipeptide and saponins such as Quil A. Particularly  
10 preferred adjuvants are, MPL (monophosphoryl lipid A) and 3D-MPL (3Deacylated monophosphoryl lipid A). A further preferred adjuvant is known as QS21. 3 DMPL can be obtained by the methods disclosed in UK patent No. 2220211, whereas QS21 can be obtained by the method disclosed in US patent No. 5,057,540. As a further exemplary alternative,  
15 the proteins can be encapsulated within microparticles such as liposomes or associated with oil in water emulsions. In yet another exemplary alternative, the proteins can be conjugated to an immuostimulating macromolecule, such as killed Bordetella or a tetanus toxoid.

20 The proteins of the present invention may be expressed by live vectors such as BCG, *Listeria* or *Salmonella* and formulated as live vaccines using such vectors.

Vaccine preparation is generally described in New Trends and  
25 Developments in Vaccines, Voller et al. (eds.), University Park Press, Baltimore, Maryland, 1978. Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877. Conjugation of proteins to macromolecules is disclosed, for example, by Likhite, US Patent 4,372,945 and Armor et al., US Patent 4,474,757.

30 Use of Quil A is disclosed by Dalsgaard et al., Acta Vet Scand. 18:349 (1977). Use of 3D-MPL is described by Ribí et al. Microbiology (1986) LEVIE et al. (Eds) American Soc. for Microbiol. Wash. D.C. pp 9-13.

35 The amount of the protein of the present invention present in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccines. Such amount will vary depending upon which specific immunogen is employed

-15-

and whether or not the vaccine is adjuvanted. Generally, it is expected that each dose will comprise 1-1000  $\mu\text{g}$  of protein, preferably 1-200  $\mu\text{g}$ . An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of antibody titres and other responses in subjects. Following an initial vaccination, subjects may receive an additional administration to enhance their immune response.

5

Example 1Characteristics of *Borrelia burgdorferi* strain ZQ1

- 5 1.1. Isolation: Midgut sample of a female *Ixodes ricinus* captured in Freiburg-Kappel from a dog on May, 31st, 1988; grown in BSK supplemented with Kanamycin and Fluoruracil; culture became positive on June, 8th 1988; aliquots were frozen at -70°C and in liquid nitrogen on June, 20th 1988.
- 10 In vitro passages usually used for experiments second and third of *B. burgdorferi* are used.
- 15 1.2 Cultivation: *Borrelia burgdorferi* strain ZQ1 is grown in Barbour-Stoenner-Kelly's medium as described for strain ZS7 (Schailble et al., J.Exp.Med. 170. 1989;). The generation time is around 30 to 40 hours (slower than strain ZS7; around 20 - 24 hours).
- 20 1.3 Plasmids: Plasmid analysis was done in parallel for strain ZQ1, ZS7, Z37, GeHo, and B31 according to the protocol of Barbour and Garon (1977): Plasmids with a molecular weight of 49, 29, 25, 20, 18 Kb and a faint band at 14 Kb are found in strain ZQ1. The 49 and 25Kb plasmids are also present in B31, Z37, GeHo and ZS7, whereas the 29 and 20Kb plasmids are only detectable in ZS7. The 18 and 14Kb plasmids seem to be unique to ZQ1. The 16Kb plasmid which is found in the other strains tested is lacking in ZQ1.
- 25 1.4 Polyacrylamide gel electrophoresis: There is a distinct pattern of proteins in strain ZQ1 when compared to other strains as revealed by polyacrylamide gel electrophoresis and silverstaining; there is no band in the range of 34kDa (Osp B); the band most probably representing Osp A has a molecular weight of around 32kDa. The corresponding band in the other strains (ZS7, Z37, GeHo, B31) migrates at 31kDa.
- 30 1.5 Western Blot analysis: monoclonal antibodies (mAbs) to *Borrelia burgdorferi* (B31, GeHo) were used, ZQ1 shows a staining pattern distinct to that of the strains ZS7 and B31; there is no staining with any of the Osp A specific mAbs with the exception of the Osp A specific mAb LA-

-17-

- 31.1 which reacts with a corresponding band of molecular mass range around 32kDa. None of the mAbs specific for Osp B of B31, ZS7 react with a protein of ZQ1. There is also no staining with the mAb LA.7 to a 20kDa antigen. mAbs specific for the 65kDa and the 70kDa antigens stain structures in ZQ1 corresponding to those in other strains.

#### Immune sera:

- The ZQ1 derived 32kDa protein (Osp A) is recognized by rabbit Immune sera (IS) to Borrelia burgdorferi B31, mouse IS to Borrelia burgdorferi B31 and mouse IS Borrelia burgdorferi ZS7. There is no staining in the range of 34kDa, IS to Borrelia burgdorferi ZQ1 reacts with Osp A, flagellin and a 20 (faint band), 50 and 65 (faint band) kDa band but not with Osp B of strain ZS7.

#### Example 2

##### 2.1 DNA preparation and PCR amplification

- B. burgdorferi cells are lysed using the detergent SDS and proteins are removed by digestion with the nonspecific serine protease, proteinase K. Cell wall debris, polysaccharides, and remaining proteins are removed by selective precipitation with CTAB (cetyltrimethylammonium bromide), and DNA is recovered from the resulting supernatant by isopropanol precipitation. Briefly, spirochetes ( $10^{10}$  cells) are washed in PBS and suspended in 9ml TE. After addition of 1ml SDS (20%) and 50 $\mu$ l proteinase K (20mg/ml), the solution was incubated for 1 hr at 37°C. Finally, 1.5ml 5M NaCl and 1.25ml CTAB (20% in 0.7 M NaCl) were added and the mixture was kept for 10 min. at 65°C. DNA was purified by phenol and chloroform/isoamylalcohol extraction and subsequently precipitated with 0.6 vol. isopropanol for 10 min. at room temperature. The pellet was washed using 70% ethanol, dried and dissolved in TE. (10mM Tris 0.1mol EDTA pH 7.4).
- Oligonucleotide primers for the polymerase chain reaction were chemically synthesized by using an Applied Biosystems (ABI) DNA synthesizer and b-cyanoethyl chemistry. The following sequences of oligonucleotides were used through out the experiments:

prOsp A1: 5'-GGGAATAGGT CTAATATTAG CC-3';  
prOsp A2: 5'-TGACCTGAATT CCAAGCTGCA-3';  
prOsp A4: 5'-GCAGTTAAAG TTCCTTCAAG AACATAGC-3'.

5

Taq DNA polymerase and PCR reagents (Perkin Elmer-Cetus, Norwalk, Conn.) were used per 50µl reaction. Each polymerase chain reaction contained 800mM deoxynucleoside triphosphate, 2U Taq DNA polymerase, 1.5mM MgCl<sub>2</sub>, and 0.5mM of each primer. Routinely, 10ng of total genomic *B. burgdorferi* DNA was amplified for 35 cycles under the following conditions: 94°C for 1.2 min; 42°C for 2 min; 72°C for 2 min. A total of 10µl per reaction was analysed on 1.5% agarose gels under standard electrophoresis conditions.

10

## 15 2.2 Southern Analysis

Total genomic *B. burgdorferi* DNA (20µg), was digested with restriction enzymes as indicated, separated by electrophoresis on a 0.8% agarose gel, and transferred to nylon membrane (Hybond-N, Amersham). Blots were hybridized at 60°C in Church-buffer (0.5M Na HPO<sub>4</sub> 17% NaDodSO<sub>4</sub>, PH 7.2) over night with an Osp A (strain ZS7) probe labelled with <sup>32</sup>P by random primer extension and washed at 55°C in 0.1x SSC/0.1% SDS.

20

## Results

25

Table 1: Genotypical and phenotypical analyses of *B. burgdorferi* strains as analysed by PCR, and Southern blotting are shown in Table I.

## 30 **Restriction Fragment length polymorphism of the plasmid encoded Osp A gene**

RFPL for Osp A using HindIII revealed at least six distinct hybridization patterns among the *B. burgdorferi* isolates tested. All of the *B. burgdorferi* strains tentatively termed as group A,A,A (for a fla, HSP60, HSP70) are characterized by two hybridization fragments of 1.2kb and 0.3kb (B31, ZS7; Osp A type I) and the majority of group B,B,B (for fla, HSP60, HSP70) strains (17/25) expressed two fragments of 0.9kb and

35

-19-

0.4kb (ZQ1; Osp A type II). *B. burgdorferi* strains 19857 and 21038 (group II,A/B,A for fla, HSP60, HSP70) exhibited two hybridizing fragments of 4kb and 0.5kb (Osp A genotype III) and all ACA-1 like strains (group II,B,A) express only on fragment of about 1.7 kb (Osp A type IV). Strains 20047 and S90 (group II,B,B) exhibit one fragment of 3kb (Osp A type V) and 1.1kb (Osp A type VI), respectively (Tables 1). DNAs isolated from members of other species of *Borrelia* did not hybridize to the Osp A-specific probe. The frequencies of the individual Osp A genotypes among *B. burgdorferi* isolates tested are: 30/55 (55%) Osp A type I, 17/55 (31%) type II, 2/55 (4%) type III, 4/55 (7%) type IV, 1/55 (2%) type V and 1/55 (2%) type VI. Note that American *B. burgdorferi* isolates analysed to this end belong to only one Osp A genotype, i.e. type I, with the exception of two strains, 19857 and 21038, which express Osp A genotype III; in contrast, among European isolates five Osp A genotypes, i.e. I, II, IV, V and VI are found. The results are summarised in table IV.

### Example 3

**Pathogenicity:** Inoculation of scid mice with  $10^8$  ZQ1 subcutaneously in the tail leads to clinical arthritis around day 50 post inoculation. Histopathological alterations in the joints investigated at day 57 were comparable to those described for scid mice infected with strain ZS7 in earlier stages of the disease. ZQ1 is therefore less pathogenic than ZS7. Spirochetes could be isolated at day 57 post inoculation from scid mice infected with ZQ1.

When 5 C57BL/6 mice were inoculated with  $10^8$  B.b. ZQ1 subcutaneously in the tail, only one mouse developed unilateral clinical arthritis at day 71 post inoculation within the entire observation period of 160 days.

### Example 4

**Crossprotection:** Reconstitution of scid mice with IS to ZQ1 (32 µg specific Ig/ml; 100µl at days 0 and 4, 200µl at days 7 and 11 and 300µl at days 15 and 19) did not protect these mice against a challenge with ZS7 given at day 0.

-20-

Example 5

- Further crossprotection studies were carried out utilising strain the Group II strain NE11H and the group I strain ZS7 Transfer of immune sera into
- 5 SCID mice as shown in Table II only protected against homologous challenge (i.e. challenge within the same group).

Example 6

- 10 Crossprotection experiments using isolates or to immune sera to either various *B. burgdorferi* isolates or to rec.Osp A of strain ZS7

**Materials and Methods**

15 Methods

- Mice (C57BL/6 or DBA/2) were inoculated with  $1 \times 10^8$  viable *B. burgdorferi* spirochetes of the strains ZS7 (IS anti-ZS7), ZQ1 (IS anti-ZQ1), NE11H (IS anti-NE11H), B31 (IS anti-B31) or a mixture containing
- 20 equal amounts of strains 20047, ZS7, NE11H, 21305, 21038, 21343, 26816, 28691 and 19535 (IS anti-cocktail) subcutaneously (sc) into the basis of the tail.

- Rec.Osp A from strain ZS7 was expressed from the pUEX1 expression
- 25 vector in *Escherichia coli*, the proteins were extracted and affinity purified using an Osp A specific mab (LA-2). Mice were primed s.c. and boosted after 10 and 20 days with 10µg rec. Osp A in adjuvant (ABM2; Sebac, Aidenbach, FRG).

- 30 Immune sera (IS) were obtained from these mice by several bleedings between week three and ten post inoculation (p.i.). SCID mice were injected intraperitoneally (i.p.) with IS either once (60 µg *B. burgdorferi* specific Ig; IS anti-ZS7, IS anti-NE11H, IS anti-cocktail, IS anti rec. Osp A) one hour before inoculation or six times for three weeks (IS anti-B31
- 35 containing 60 µg/ml specific Ig; IS anti-ZQ1 containing 32 µg/ml) with increasing amounts of antibodies (100µl day 1 and 4, 200µl day 7 and 11,300 µl day 14 and 17 p.i.). SCID mice were inoculated with  $1 \times 10^8$  viable *B. burgdorferi* spirochetes of the strains ZS7, SH-2-82-P5, 21038,

-21-

NE11H or 20047 s.c. into the base of the tail one hour after the (first) transfer of serum.

Results:

5

Immune sera to subgroup I strains ZS7 and B31 protect against homologous challenge, only partially against heterologous challenge with strain 20047 (subgroup V) and not at all against heterologous challenge with subgroup II strain NE11H. Immune sera to group II strain NE11H only partially protects against heterologous challenge with strain 20047 (subgroup V) and not at all against heterologous challenge with strain ZS7 (subgroup I).

10

Pretreatment of SCID mice with monospecific anti-rec. Osp A (ZS7) IS resulted in considerably protection against development of disease when challenged with either strain ZS7 (2/6) or SH-2-82-P5 (2/3; flaA, OspA/OspB I) and in partial protection when challenged with 20047 (1/3; flaB, Osp A/Osp B V) or 21038 (2/3; fla B, Osp A/Osp B III; Table 2). The clinical arthritis observed in the majority of these mice under these conditions was only mild. In contrast, SCID mice treated with the same IS to rec. Osp A but subsequently challenged with NE11H (fla B, Osp A/Osp B II) were not protected at all. All of these mice developed clinical arthritis (3/3) comparable to that seen in infected and otherwise untreated SCID mice (Table 3).

20

25



Example 7Construction of expression plasmids encoding NS1-OspA (ZS7) fusion

- 5 Figure 1 outlines the construction of the three plasmids pOA-10, POA-12 and pOA-18.
- Briefly, pOA-10 was created in three steps. First, a 1.1 kb NspH 1 fragment of pZS7-2 (Dr Wallich) containing the truncated OspA 'gene was ligated, after trimming of the protruding ends, with pUC19 linearized
- 10 with Sma 1. In the resulting plasmid, named pOA-5, the OspA gene is flanked by various unique restriction sites,(in 5' end these are Hind III, Sph I, Pst I Sal I, Xba and BamH I and in the 3' end Sac 1 and EcoR I) making easier its subcloning. Secondly, pOA-5 was linearized by Xba I and ligated with pMG42 linearized by BamH I, after fill-in of the ends of
- 15 both fragments to create pOA-9. The portion of this plasmid derived from pUC19 has subsequently been removed by an Nco I digestion, generating pOA-10. In this plasmid, the truncated OspA gene is in frame with the first codon of NS1. The junction at the 5' end of the insert has been
- 20 verified by sequencing. To create pOA-12 and pOA-18, we generated Nco I and Xba I restriction sites respectively at the 5' and the 3' ends of the DNA fragment coding for the truncated OspA gene using polymerase chain reaction (PCR). After digestion with Nco I and Xba I, this fragment was ligated with pMG42 or TCM67 digested with the same enzymes to obtain respectively pOA-12 and pOA-18. The first plasmid pOA-12,
- 25 expresses the truncated OspA gene fused to the first 42 amino acids, the latter, pOA-18, to the first 81 amino acids of NS1. The insert of both plasmids have been sequenced to verify that no amino acid substitution has been introduced in OspA gene during the PCR.
- 30 All three plasmids have been transformed into *E. coli* AR120 which allows the induction the  $\lambda$ pL promoter by the addition of nalidixic acid. Additionally all three plasmids have been transformed into *E. coli* AR58 which allows induction of the  $\lambda$ pL promoter by an increase in
- 35 temperature.

Example 8Expression of recombinant trOspA from pOA-10, pOA-12 and POA-18

- 5 Fifty ml cultures (LB medium supplemented with the appropriate antibiotics) of the three strains AR120 (pOA-10), AR120 (pOA-12) and AR120 (pOA-18) were inoculated with 0.25 ml of an overnight preculture and grown at 37°C. At an  $A_{620}$  of about 0.3, the cells contained in 5 ml of culture have been harvested and resuspended in SDS sample buffer.
- 10 Nalidixic acid has been added to the remaining culture to a final concentration of 60mg/ml. Three hours later the cells from the 5ml cultures were harvested and resuspended in SDS sample buffer.

- 15 In parallel to the three AR120 strains, JM109 and JM109 (pZS7-2) strains have been grown in 50 ml of LB medium up to an optical density (620 nm) of 0.6 and treated as described above. In JM109(pZS7-2), the native form of OspA is expressed constitutively from its own promoter. All samples were submitted to SDS-PAGE and Western blot analysis using a pool of the mAbs LA-2, LA-26, and LA-31 to detect the OspA derivatives.

- 20 Analysis of samples corresponding to uninduced and induced cells (in the case of pZS7-2, host without and with the plasmid), were carried out. The amount of material loaded into each well was standardized on the basis of the optical density of the cultures (0.05  $A_{620}$ /well). On the stained gel, in
- 25 the lane induced of JM109(pZS7-2) and of induced AR120(pOA-18), new bands appear which migrations are consistent with the expected size of the products to be expressed (respectively native OspA and trOspA fused to 81 amino acids of NS1). Native OspA appears as a doublet and it represents a few percents of the total protein content of the bacteria. The
- 30 product of pOA-18 is expressed at higher level than the native protein. The Western blot analysis confirms that these new bands correspond to OspA derivatives. The blot further shows that AR120(pOA-12) expresses trOspA fused with 42 amino acids of NS1 but at a lower level. The expression of trOspA from pOA-10 appears as a faint band, corresponding
- 35 to the expected size of trOspA and was observed on Western blot.

Similar results have been observed when the same plasmids are introduced into E. coli AR58 and the expression of the Osp A gene

-24-

derivative induced by an increase of temperature. Also we have obtained similar NS1<sub>1-81</sub> Osp A ZS7 fusion protein utilising a kanamycin resistance encoding plasmid.

## 5 Example 9

### Comparison of cells fractionation of AR120(pOA18) and JM109(pZS7-2)

10 E. coli strain AR120(pOA18) was grown, and the expression of the Osp A genes induced as described above.

15 E. coli strain JM109(pZS7-2), in which the native gene of Osp A is expressed constitutively from its own promoter, has been grown in 50 ml of LB medium supplemented with 100 µg/ml of ampicillin up to an optical density (620 nm) of 0.6.

20 The cells of both strains were then treated as follows. The cells contained in 5 ml have been harvested by centrifugation and resuspended in SDS-PAGE sample buffer (sample T3). The cells in the remaining 40 ml were harvested, washed twice in buffer A (Tris-HCl 50 mM, pH 7.5) and disrupted in the same buffer by two passages at 10,000 lb/in<sup>2</sup> through a French pressure cell. Cell homogenates were fractionated by centrifugation. The supernatants were stored at -20°C (samples D3). The pellets were washed twice in buffer A and solubilized overnight in SDS-  
25 PAGE sample buffer at room temperature (sample C3). Results of the cell fractionation of the 2 expressing strains were analyzed by SDS-PAGE and Western blot was conducted as above except that only 0.025 A<sub>620</sub> of "cell-equivalent" was loaded into each well. The native OspA, expressed in JM109(pZS7-2), is found in both the soluble (S3) and the insoluble (C3)  
30 cellular fractions. The lower band of the doublet is found preferentially in the insoluble fraction, while the upper band is almost entirely in the soluble fraction. The high molecular weight protein that appear only in the C3 lane of JM109(pZS7-2) is not recognized by the anti-OspA antibodies and is thus probably not OspA-derived. On the other hand the  
35 product of pOA-18 is found almost exclusively in the soluble fraction.

-25-

Example 10Reactivity of NSI-OspA from pOA-18 with 4 anti-OspA mAbs

5 Crude cell extracts pf AR120 (pOA18) and JM109(pZS7-2) were prepared as described above. They were submitted to SDS-PAGE in four series of side by side wells (AR120(pOA-18) and JM109(pZS7-2)) and their proteins transferred onto a nitrocellulose membrane. The membrane were then cut in four fragments each containing a pair of side by side lanes. Each  
10 fragment was submitted to a Western blot analysis with a different anti-Osp A mAb. The results indicate that the NSI- Osp A encoded by pOA-18 and the native Osp A react with all four monoclonals in a similar fashion.

Example 11

15 In an analogous fashion to the construction of pOA-18, plasmids were constructed to express NS1-OspA fusions proteins from other B. burgdorferi strain. In particular the following fusion from the following strains were made and tested as noted above: NS1-OspA (ZQ1); NS1-  
20 OspA (AcaI) and NS1-OspA (19857).

Summary and conclusions

• A recombinant truncated form of OspA, lacking the 16 first amino  
25 acids of the native protein and fused to the first 81 amino acids of NS1, is expressed at a higher level than the native OspA. It is more soluble than the native protein and retains the reactivity with those monoclonal antibodies binding to the respective native Osp A is immunogenic when applied with adjuvants and has the ability to protect animals from  
30 challenge with virulent B. burgdorferi.

-26-

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(i) APPLICANT: SIMON, MARKUS  
KRAMER, MICHAEL  
SCHAIBLE, ULRICH  
WALLICH, REINHARD  
LOBET, YVES

(ii) TITLE OF INVENTION: VACCINES

(iii) NUMBER OF SEQUENCES: 12

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: MARCUS DALTON SMITHKLINE BEECHAM  
(B) STREET: GREAT BURGH YEW TREE BOTTOM ROAD  
(C) CITY: EPSOM  
(D) STATE: SURREY  
(E) COUNTRY: UNITED KINGDOM  
(F) ZIP: KT18 5XQ

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: DALTON, MARCUS JW  
(B) REGISTRATION NUMBER: 22885  
(C) REFERENCE/DOCKET NUMBER: B45008

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 0737 364000  
(B) TELEFAX: 0737 364006  
(C) TELEX: 911415

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 825 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: BORRELIA BURGDORFERI  
(B) STRAIN: ZQ1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAAAAAAT ATTTATTGGG AATAGGTCTA ATATTAGCCT TAATAGCATG TAAGCAAAAT  
60

GTTAGCAGCC TTGATGAAAA AAATAGCGTT TCAGTAGATT TACCTGGTGG AATGAAAGTT  
120

-27-

CTTGTAAGTA AAGAAAAAGA CAAAGATGGT AAATACAGTC TAGAGGCAAC AGTAGACAAG  
180

CTTGAGCTTA AAGGAACTTC TGATAAAAAC AACGGTTCTG GAACACTTGA AGGTGAAAAA  
240

ACTGACAAAA GTAAAGTAAA ATTAACAATT GCTGAGGATC TAAGTAAAAC CACATTTGAA  
300

ATTTTCAAAG AAGATGGCAA AACATTAGTA TCAAAAAAAG TAACCCTTAA AGACAAGTCA  
360

TCAACAGAAG AAAAATTCAA CGAAAAGGGT GAAATATCTG AAAAAACAAT AGTAAGAGCA  
420

AATGGAACCA GACTTGAATA CACAGACATA AAAAGCGATG GATCCGAAA AGCTAAAGAA  
480

GTTTTAAAAG ACTTTACTCT TGAAGGAACT CTAGCTGCTG ACGGCAAAAC AACATTGAAA  
540

GTTACAGAAG GCACTGTTGT TTTAAGCAAG AACATTTTAA AATCCGGAGA AATAACAGTT  
600

GCACTTGATG ACTCTGACAC TACTCAGGCT ACTAAAAAA CTGGAATG GGATTCAAAG  
660

ACTTCCACTT TAACAATTAG TGTGAATAGC CAAAAAACCA AAACCTTGT ATTCACAAAA  
720

GAAGACACAA TAACAGTACA AAAATACGAC TCAGCAGGCA CCAATCTAGA AGGCAAAGCA  
780

GTCGAAATTA CAACACTTAA AGAACTTAA GACGCTTTAA AATAA  
825

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 274 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: ZQ1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Lys	Lys	Tyr	Leu	Leu	Gly	Ile	Gly	Leu	Ile	Leu	Ala	Leu	Ile	Ala
1					5				10					15	
Cys	Lys	Gln	Asn	Val	Ser	Ser	Leu	Asp	Glu	Lys	Asn	Ser	Val	Ser	Val
			20					25					30		
Asp	Leu	Pro	Gly	Gly	Met	Lys	Val	Leu	Val	Ser	Lys	Glu	Lys	Asp	Lys
		35					40					45			
Asp	Gly	Lys	Tyr	Ser	Leu	Glu	Ala	Thr	Val	Asp	Lys	Leu	Glu	Leu	Lys
	50					55					60				
Gly	Thr	Ser	Asp	Lys	Asn	Asn	Gly	Ser	Gly	Thr	Leu	Glu	Gly	Glu	Lys
65					70					75				80	

-28-

Thr Asp Lys Ser Lys Val Lys Leu Thr Ile Ala Glu Asp Leu Ser Lys  
 85 90 95  
 Thr Thr Phe Glu Ile Phe Lys Glu Asp Gly Lys Thr Leu Val Ser Lys  
 100 105 110  
 Lys Val Thr Leu Lys Asp Lys Ser Ser Thr Glu Glu Lys Phe Asn Glu  
 115 120 125  
 Lys Gly Glu Ile Ser Glu Lys Thr Ile Val Arg Ala Asn Gly Thr Arg  
 130 135 140  
 Leu Glu Tyr Thr Asp Ile Lys Ser Asp Gly Ser Gly Lys Ala Lys Glu  
 145 150 155 160  
 Val Leu Lys Asp Phe Thr Leu Glu Gly Thr Leu Ala Ala Asp Gly Lys  
 165 170 175  
 Thr Thr Leu Lys Val Thr Glu Gly Thr Val Val Leu Ser Lys Asn Ile  
 180 185 190  
 Leu Lys Ser Gly Glu Ile Thr Val Ala Leu Asp Asp Ser Asp Thr Thr  
 195 200 205  
 Gln Ala Thr Lys Lys Thr Gly Lys Trp Asp Ser Lys Thr Ser Thr Leu  
 210 215 220  
 Thr Ile Ser Val Asn Ser Gln Lys Thr Lys Asn Leu Val Phe Thr Lys  
 225 230 235 240  
 Glu Asp Thr Ile Thr Val Gln Lys Tyr Asp Ser Ala Gly Thr Asn Leu  
 245 250 255  
 Glu Gly Lys Ala Val Glu Ile Thr Thr Leu Lys Glu Leu Lys Asp Ala  
 260 265 270  
 Leu Lys

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 819 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: 19857

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGAAAAAAT ATTTATTGGG AATAGGTCTA ATATTAGCCT TAATAGCATG TAAGCAAAAT  
60

GTTAGCAGCC TTGACGAGAA AACAGCGTT TCAGTAGATG TACCTGGTGG AATGAAAGTT  
120

CTTGTAAGCA AAGAAAAAAA CAAAGACGGC AAGTACGATC TAATGGCAAC AGTGGACAAC  
180

GTTGATCTTA AAGGAACTTC TGACAAAAAC AATGGATCTG GAATACTTGA AGGCGTAAAA  
240

-29-

GCTGATAAAA GTAAAGTAAA ATTAACAGTT GCTGACGATC TAAGCAAAAC CACACTTGAA  
300

GTTTTAAAAG AAGATGGTAC AGTAGTGTCAG AAAAAGTAA CTTCCAAAGA CAAGTCAACA  
360

ACAGAAGCAA AATTCAACGA AAAAGGTGAA TTGTCTGAAA AAACAATGAC AAGAGCAAAC  
420

GGAAGTACTC TTGAATCTC ACAAATGACA AATGAAGACA ATGCTGCAAA AGCAGTAGAA  
480

ACTCTTAAAA ACGGCATTAA GTTTGAAGGA AATCTCGCAA GTGGAAAAAC AGCAGTGGAA  
540

ATTAAAGAAG GCACTGTTAC TCTAAAAAGA GAAATTGATA AAAATGGAAA AGTAACCGTC  
600

TCTTTAAATG ACACTGCATC TGGTTCTAAA AAAACAGCTT CCTGGCAAGA AAGTACTAGC  
660

ACCTTAACAA TTAGTGCAAA CAGCAAAAAA ACTAAAGATC TAGTGTTCTT AACAAACGGT  
720

ACAATTACAG TACAAAATTA TGAATCAGCT GGCATAAAC TTGAAGGATC AGCAGCTGAA  
780

ATTAAAAAAC TCGATGAACT TAAAAACGCT TTAAGATAA  
819

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 272 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: BORRELIA BURGDORFERI  
 (B) STRAIN: 19857

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Lys	Lys	Tyr	Leu	Leu	Gly	Ile	Gly	Leu	Ile	Leu	Ala	Leu	Ile	Ala
1				5					10					15	
Cys	Lys	Gln	Asn	Val	Ser	Ser	Leu	Asp	Glu	Lys	Asn	Ser	Val	Ser	Val
			20					25					30		
Asp	Val	Pro	Gly	Gly	Met	Lys	Val	Leu	Val	Ser	Lys	Glu	Lys	Asn	Lys
		35					40					45			
Asp	Gly	Lys	Tyr	Asp	Leu	Met	Ala	Thr	Val	Asp	Asn	Val	Asp	Leu	Lys
	50					55					60				
Gly	Thr	Ser	Asp	Lys	Asn	Asn	Gly	Ser	Gly	Ile	Leu	Glu	Gly	Val	Lys
65					70					75				80	
Ala	Asp	Lys	Ser	Lys	Val	Lys	Leu	Thr	Val	Ala	Asp	Asp	Leu	Ser	Lys
				85					90					95	
Thr	Thr	Leu	Glu	Val	Leu	Lys	Glu	Asp	Gly	Thr	Val	Val	Ser	Arg	Lys
				100				105						110	



-30-

Val Thr Ser Lys Asp Lys Ser Thr Thr Glu Ala Lys Phe Asn Glu Lys  
115 120 125

Gly Glu Leu Ser Glu Lys Thr Met Thr Arg Ala Asn Gly Thr Thr Leu  
130 135 140

Glu Tyr Ser Gln Met Thr Asn Glu Asp Asn Ala Ala Lys Ala Val Glu  
145 150 155 160

Thr Leu Lys Asn Gly Ile Lys Phe Glu Gly Asn Leu Ala Ser Gly Lys  
165 170 175

Thr Ala Val Glu Ile Lys Glu Gly Thr Val Thr Leu Lys Arg Glu Ile  
180 185 190

Asp Lys Asn Gly Lys Val Thr Val Ser Leu Asn Asp Thr Ala Ser Gly  
195 200 205

Ser Lys Lys Thr Ala Ser Trp Gln Glu Ser Thr Ser Thr Leu Thr Ile  
210 215 220

Ser Ala Asn Ser Lys Lys Thr Lys Asp Leu Val Phe Leu Thr Asn Gly  
225 230 235 240

Thr Ile Thr Val Gln Asn Tyr Asp Ser Ala Gly Thr Lys Leu Glu Gly  
245 250 255

Ser Ala Ala Glu Ile Lys Lys Leu Asp Glu Leu Lys Asn Ala Leu Arg  
260 265 270

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1020 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: zql

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: NS1-ZQ1

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATGGATCCAA ACACTGTGTC AAGCTTTCAG GTAGATTCCT TTCTTTGGCA TGTCCGCAAA  
60


CGAGTTGCAG ACCAAGAACT AGGTGATGCC CCATTCCTTG ATCGGCTTCG CCGAGATCAG  
120

AAATCCCTAA GAGGAAGGGG CAGCACTCTT GGTCTGGACA TCGAGACAGC CACACGTGCT  
180

GGAAAAGCAGA TAGTGGAGCG GATTCTGAAA GAAGAATCCG ATGAGGCACT TAAAATGACC  
240

ATGGGAAAGC AAAATGTTAG CAGCCTTGAT GAAAAAATA GCGTTTCAGT AGATTACCT  
300

GGTGGAATGA AAGTTCTTGT AAGTAAAGAA AAAGACAAAG ATGGTAAATA CAGTCTAGAG  
360



-31-

GCAACAGTAG ACAAGCTTGA GCTTAAAGGA ACTTCTGATA AAAACAACGG TTCTGGAACA  
420

CTTGAAGGTG AAAAACTGA CAAAAGTAAA GTAAAATTAA CAATTGCTGA GGATCTAAGT  
480

AAAACCACAT TTGAAATTTT CAAAGAAGAT GGCAAAACAT TAGTATCAAA AAAAGTAACC  
540

CTTAAAGACA AGTCATCAAC AGAAGAAAAA TTCAACGAAA AGGGTGAAAT ATCTGAAAAA  
600

ACAATAGTAA GAGCAAATGG AACCAGACTT GAATACACAG ACATAAAAAG CGATGGATCC  
660

GGAAAAGCTA AAGAAGTTTT AAAAGACTTT ACTCTTGAAG GAACTCTAGC TGCTGACGGC  
720

AAAACAACAT TGAAAGTTAC AGAAGGCACT GTTGTTTTAA GCAAGAACAT TTTAAATCC  
780

GGAGAAATAA CAGTTGCACT TGATGACTCT GACACTACTC AGGCTACTAA AAAAACTGGA  
840

AAATGGGATT CAAAGACTTC CACTTTAACA ATTAGTGTGA ATAGCCAAAA AACCAAAAC  
900

CTTGTATTCA CAAAAGAAGA CACAATAACA GTACAAAAAT ACGACTCAGC AGGCACCAAT  
960

CTAGAAGGCA AAGCAGTCGA AATTACAACA CTTAAGAAC TTAAAGACGC TTTAAATAA  
1020

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: ZQ1

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: NS1-ZQ1

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Asp	Pro	Asn	Thr	Val	Ser	Ser	Phe	Gln	Val	Asp	Ser	Phe	Leu	Trp
1				5					10					15	
His	Val	Arg	Lys	Arg	Val	Ala	Asp	Gln	Glu	Leu	Gly	Asp	Ala	Pro	Phe
			20					25					30		
Leu	Asp	Arg	Leu	Arg	Arg	Asp	Gln	Lys	Ser	Leu	Arg	Gly	Arg	Gly	Ser
		35					40					45			
Thr	Leu	Gly	Leu	Asp	Ile	Glu	Thr	Ala	Thr	Arg	Ala	Gly	Lys	Gln	Ile
		50				55						60			

-32-

Val Glu Arg Ile Leu Lys Glu Glu Ser Asp Glu Ala Leu Lys Met Thr  
 65 70 75 80  
 Met Gly Lys Gln Asn Val Ser Ser Leu Asp Glu Lys Asn Ser Val Ser  
 85 90 95  
 Val Asp Leu Pro Gly Gly Met Lys Val Leu Val Ser Lys Glu Lys Asp  
 100 105 110  
 Lys Asp Gly Lys Tyr Ser Leu Glu Ala Thr Val Asp Lys Leu Glu Leu  
 115 120 125  
 Lys Gly Thr Ser Asp Lys Asn Asn Gly Ser Gly Thr Leu Glu Gly Glu  
 130 135 140  
 Lys Thr Asp Lys Ser Lys Val Lys Leu Thr Ile Ala Glu Asp Leu Ser  
 145 150 155 160  
 Lys Thr Thr Phe Glu Ile Phe Lys Glu Asp Gly Lys Thr Leu Val Ser  
 165 170 175  
 Lys Lys Val Thr Leu Lys Asp Lys Ser Ser Thr Glu Glu Lys Phe Asn  
 180 185 190  
 Glu Lys Gly Glu Ile Ser Glu Lys Thr Ile Val Arg Ala Asn Gly Thr  
 195 200 205  
 Arg Leu Glu Tyr Thr Asp Ile Lys Ser Asp Gly Ser Gly Lys Ala Lys  
 210 215 220  
 Glu Val Leu Lys Asp Phe Thr Leu Glu Gly Thr Leu Ala Ala Asp Gly  
 225 230 235 240  
 Lys Thr Thr Leu Lys Val Thr Glu Gly Thr Val Val Leu Ser Lys Asn  
 245 250 255  
 Ile Leu Lys Ser Gly Glu Ile Thr Val Ala Leu Asp Asp Ser Asp Thr  
 260 265 270  
 Thr Gln Ala Thr Lys Lys Thr Gly Lys Trp Asp Ser Lys Thr Ser Thr  
 275 280 285  
 Leu Thr Ile Ser Val Asn Ser Gln Lys Thr Lys Asn Leu Val Phe Thr  
 290 295 300  
 Lys Glu Asp Thr Ile Thr Val Gln Lys Tyr Asp Ser Ala Gly Thr Asn  
 305 310 315 320  
 Leu Glu Gly Lys Ala Val Glu Ile Thr Thr Leu Lys Glu Leu Lys Asp  
 325 330 335  
 Ala Leu Lys

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1014 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: 19857

-33-

(vii) IMMEDIATE SOURCE:  
(B) CLONE: NS1-19857

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGGATCCAA ACACTGTGTC AAGCTTTCAG GTAGATTCCT TTCTTTGGCA TGTCCGCAAA  
60  
CGAGTTGCAG ACCAAGAAGT AGGTGATGCC CCATTCCTTG ATCGGCTTCG CCGAGATCAG  
120  
AAATCCCTAA GAGGAAGGGG CAGCACTCTT GGTCTGGACA TCGAGACAGC CACACGTGCT  
180  
GGAAAGCAGA TAGTGGAGCG GATTCTGAAA GAAGAATCCG ATGAGGCACT TAAATGACC  
240  
ATGGGAAAGC AAAATGTTAG CAGCCTTGAC GAGAAAAACA GCGTTTCAGT AGATGTACCT  
300  
GGTGGAAATGA AAGTTCTTGT AAGCAAAGAA AAAAACAAG ACGGCAAGTA CGATCTAATG  
360  
GCAACAGTGG ACAACGTTGA TCTTAAAGGA ACTTCTGACA AAAACAATGG ATCTGGAATA  
420  
CTTGAAGGCG TAAAAGCTGA TAAAAGTAAA GTAAATTAA CAGTTGCTGA CGATCTAAGC  
480  
AAAACACAC TTGAAGTTT AAAAGAAGAT GTACAGTAG TGCAAGAAA AGTAACCTCC  
540  
AAAGACAAGT CAACAACAGA AGCAAATTC AACGAAAAG GTGAATTGTC TGAAAAACA  
600  
ATGACAAGAG CAAACGGAAC TACTCTTGAA TACTCACAAA TGACAAATGA AGACAATGCT  
660  
GCAAAAGCAG TAGAACTCT TAAAACGGC ATTAAGTTTG AAGGAAATCT CGCAAGTGGA  
720  
AAAACAGCAG TGGAAATTAA AGAAGGCACT GTTACTCTAA AAAGAGAAAT TGATAAAAT  
780  
GGAAAAGTAA CCGTCTCTTT AAATGACACT GCATCTGGTT CTAAAAAAC AGCTTCCTGG  
840  
CAAGAAAGTA CTAGCACCTT AACAATTAGT GCAAACAGCA AAAAACTAA AGATCTAGTG  
900  
TTCCTAACAA ACGGTACAAT TACAGTACAA AATTATGACT CAGCTGGCAC TAACTTGAA  
960  
GGATCAGCAG CTGAAATTAA AAAACTCGAT GAACTTAAAA ACGCTTTAAA ATAA  
1014

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 337 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

-34-

(A) ORGANISM: BORRELIA BURGDORFERI  
(B) STRAIN: 19857

(vii) IMMEDIATE SOURCE:  
(B) CLONE: NS1-19857

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Asp	Pro	Asn	Thr	Val	Ser	Ser	Phe	Gln	Val	Asp	Ser	Phe	Leu	Trp	1	5	10	15
His	Val	Arg	Lys	Arg	Val	Ala	Asp	Gln	Glu	Leu	Gly	Asp	Ala	Pro	Phe	20	25	30	
Leu	Asp	Arg	Leu	Arg	Arg	Asp	Gln	Lys	Ser	Leu	Arg	Gly	Arg	Gly	Ser	35	40	45	
Thr	Leu	Gly	Leu	Asp	Ile	Glu	Thr	Ala	Thr	Arg	Ala	Gly	Lys	Gln	Ile	50	55	60	
Val	Glu	Arg	Ile	Leu	Lys	Glu	Glu	Ser	Asp	Glu	Ala	Leu	Lys	Met	Thr	65	70	75	80
Met	Gly	Lys	Gln	Asn	Val	Ser	Ser	Leu	Asp	Glu	Lys	Asn	Ser	Val	Ser	85	90	95	
Val	Asp	Val	Pro	Gly	Gly	Met	Lys	Val	Leu	Val	Ser	Lys	Glu	Lys	Asn	100	105	110	
Lys	Asp	Gly	Lys	Tyr	Asp	Leu	Met	Ala	Thr	Val	Asp	Asn	Val	Asp	Leu	115	120	125	
Lys	Gly	Thr	Ser	Asp	Lys	Asn	Asn	Gly	Ser	Gly	Ile	Leu	Glu	Gly	Val	130	135	140	
Lys	Ala	Asp	Lys	Ser	Lys	Val	Lys	Leu	Thr	Val	Ala	Asp	Asp	Leu	Ser	145	150	155	160
Lys	Thr	Thr	Leu	Glu	Val	Leu	Lys	Glu	Asp	Gly	Thr	Val	Val	Ser	Arg	165	170	175	
Lys	Val	Thr	Ser	Lys	Asp	Lys	Ser	Thr	Thr	Glu	Ala	Lys	Phe	Asn	Glu	180	185	190	
Lys	Gly	Glu	Leu	Ser	Glu	Lys	Thr	Met	Thr	Arg	Ala	Asn	Gly	Thr	Thr	195	200	205	
Leu	Glu	Tyr	Ser	Gln	Met	Thr	Asn	Glu	Asp	Asn	Ala	Ala	Lys	Ala	Val	210	215	220	
Glu	Thr	Leu	Lys	Asn	Gly	Ile	Lys	Phe	Glu	Gly	Asn	Leu	Ala	Ser	Gly	225	230	235	240
Lys	Thr	Ala	Val	Glu	Ile	Lys	Glu	Gly	Thr	Val	Thr	Leu	Lys	Arg	Glu	245	250	255	
Ile	Asp	Lys	Asn	Gly	Lys	Val	Thr	Val	Ser	Leu	Asn	Asp	Thr	Ala	Ser	260	265	270	
Gly	Ser	Lys	Lys	Thr	Ala	Ser	Trp	Gln	Glu	Ser	Thr	Ser	Thr	Leu	Thr	275	280	285	
Ile	Ser	Ala	Asn	Ser	Lys	Lys	Thr	Lys	Asp	Leu	Val	Phe	Leu	Thr	Asn	290	295	300	
Gly	Thr	Ile	Thr	Val	Gln	Asn	Tyr	Asp	Ser	Ala	Gly	Thr	Lys	Leu	Glu	305	310	315	320

Gly Ser Ala Ala Glu Ile Lys Lys Leu Asp Glu Leu Lys Asn Ala Leu  
325 330 335

**Lys**

(2) INFORMATION FOR SEQ ID NO:9:

**(i) SEQUENCE CHARACTERISTICS:**

- (A) LENGTH: 1017 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)

(v1) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI  
(B) STRAIN: ACA1

(vii) IMMEDIATE SOURCE:

- (B) CLONE: NS1-ACA1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGGATCCAA ACACTGTGTC AAGCTTTCAG GTAGATTCTT TTCTTTGGCA TGTCCGCAAA  
60

CGAGTTGCAG ACCAAGAACT AGGTGATGCC CCATTCCTTG ATCGGCTTCG CCGAGATCAG  
120

AAATCCCTAA GAGGAAGGGG CAGCACTCTT GGTCTGGACA TCGAGACAGC CACACGTGCT  
180

GGAAAGCAGA TAGTGGAGCG GATTCTGAAA GAAGAATCCG ATGAGGCACT TAAAATGACC  
240

ATGGGAAAGC AAAATGTTAG CAGCCTTGAT GAAAAAACA GCGCTTCAGT AGATTTCCT  
300

GGTGAGATGA AAGTTCTTGT AAGTAAAGAA AAAGACAAAG ACGGTAAGTA CAGTCTAAAG  
360

GCAACAGTAG ACAAGATTGA GCTAAAAGGA ACTTCTGATA AAGACAATGG TTCTGGAGTG  
420

CTTGAAGGTA CAAAAGATGA CAAAAGTAAA GCAAAATTAA CAATTGCTGA CGATCTAAGT  
480

AAAACCACAT TCGAACTTTT CAAAGAAGAT GGCAAAACAT TAGTGTCAAG AAAAGTAAGT  
540

TCTAAAGACA AAACATCAAC AGATGAAATG TTCAATGAAA AAGGTGAATT GTCTGCAAAA  
600

ACCATGACAA GAGAAAATGG AACCAAACCTT GAATATACAG AAATGAAAAG CGATGGAACC  
660

GGAAAAGCTA AAGAAGTTTT AAAAACTTT ACTCTTGAAG GAAAAGTAGC TAATGATAAA  
720

GTAACATTGG AAGTAAAAGA AGGAACCGTT ACTTTAAGTA AGGAAATTGC AAAATCTGGA  
780

GAAGTAACAG TTGCTCTTAA TGACACTAAC ACTACTCAGG CTACTAAAAA AACTGGCGCA  
840

-36-

TGGGATTCAA AAACCTTCTAC TTTAACAATT AGTGTTAACA GCAAAAAAAC TACACAACCTT  
900

GTGTTTACTA AACAAGACAC AATAACTGTA CAAAAATACG ACTCCGCAGG TACCAATTTA  
960

GAAGGCACAG CAGTCGAAAT TAAACACTT GATGAACTTA AAAACGCTTT AAAATAA  
1017

[2] INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 338 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORELLIA BURGDORFERI
- (B) STRAIN: ACAI

(vii) IMMEDIATE SOURCE:

- (B) CLONE: NS1-ACAI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met	Asp	Pro	Asn	Thr	Val	Ser	Ser	Phe	Gln	Val	Asp	Ser	Phe	Leu	Trp
1				5					10					15	
His	Val	Arg	Lys	Arg	Val	Ala	Asp	Gln	Glu	Leu	Gly	Asp	Ala	Pro	Phe
			20					25					30		
Leu	Asp	Arg	Leu	Arg	Arg	Asp	Gln	Lys	Ser	Leu	Arg	Gly	Arg	Gly	Ser
		35					40					45			
Thr	Leu	Gly	Leu	Asp	Ile	Glu	Thr	Ala	Thr	Arg	Ala	Gly	Lys	Gln	Ile
	50					55					60				
Val	Glu	Arg	Ile	Leu	Lys	Glu	Glu	Ser	Asp	Glu	Ala	Leu	Lys	Met	Thr
	65				70					75				80	
Met	Gly	Lys	Gln	Asn	Val	Ser	Ser	Leu	Asp	Glu	Lys	Asn	Ser	Ala	Ser
				85					90					95	
Val	Asp	Leu	Pro	Gly	Glu	Met	Lys	Val	Leu	Val	Ser	Lys	Glu	Lys	Asp
			100					105					110		
Lys	Asp	Gly	Lys	Tyr	Ser	Leu	Lys	Ala	Thr	Val	Asp	Lys	Ile	Glu	Leu
		115					120					125			
Lys	Gly	Thr	Ser	Asp	Lys	Asp	Asn	Gly	Ser	Gly	Val	Leu	Glu	Gly	Thr
	130					135					140				
Lys	Asp	Asp	Lys	Ser	Lys	Ala	Lys	Leu	Thr	Ile	Ala	Asp	Asp	Leu	Ser
	145				150					155				160	
Lys	Thr	Thr	Phe	Glu	Leu	Phe	Lys	Glu	Asp	Gly	Lys	Thr	Leu	Val	Ser
			165						170					175	
Arg	Lys	Val	Ser	Ser	Lys	Asp	Lys	Thr	Ser	Thr	Asp	Glu	Met	Phe	Asn
			180					185					190		
Glu	Lys	Gly	Glu	Leu	Ser	Ala	Lys	Thr	Met	Thr	Arg	Glu	Asn	Gly	Thr
		195					200					205			

-37-

Lys Leu Glu Tyr Thr Glu Met Lys Ser Asp Gly Thr Gly Lys Ala Lys  
 210 215 220  
 Glu Val Leu Lys Asn Phe Thr Leu Glu Gly Lys Val Ala Asn Asp Lys  
 225 230 235 240  
 Val Thr Leu Glu Val Lys Glu Gly Thr Val Thr Leu Ser Lys Glu Ile  
 245 250 255  
 Ala Lys Ser Gly Glu Val Thr Val Ala Leu Asn Asp Thr Asn Thr Thr  
 260 265 270  
 Gln Ala Thr Lys Lys Thr Gly Ala Trp Asp Ser Lys Thr Ser Thr Leu  
 275 280 285  
 Thr Ile Ser Val Asn Ser Lys Lys Thr Thr Gln Leu Val Phe Thr Lys  
 290 295 300  
 Gln Asp Thr Ile Thr Val Gln Lys Tyr Asp Ser Ala Gly Thr Asn Leu  
 305 310 315 320  
 Glu Gly Thr Ala Val Glu Ile Lys Thr Leu Asp Glu Leu Lys Asn Ala  
 325 330 335  
 Leu Lys

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1017 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: BORRELIA BURGDORFERI
- (B) STRAIN: ZS7

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: NS1-ZS7

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATGGATCCAA ACACTGTGTC AAGCTTTCAG GTAGATTCCT TTCTTTGGCA TGTCCGCAAA  
60

CGAGTTGCAG ACCAAGAACT AGGTGATGCC CCATTCCTTG ATCGGCTTCG CCGAGATCAG  
120

AAATCCCTAA GAGGAAGGGG CAGCACTCTT GGTCTGGACA TCGAGACAGC CACACGTGCT  
180

GGAAAGCAGA TAGTGGAGCG GATTCTGAAA GAAGAATCCG ATGAGGCACT TAAAATGACC  
240

ATGGGAAAGC AAAATGTTAG CAGCCTTGAC GAGAAAAACA GCGTTTCAGT AGATTTCCTT  
300

GGTGAAATGA ACGTTCTTGT AAGCAAAGAA AAAAACAAG ACGGCAAGTA CGATCTAATT  
360

GCAACAGTAG ACAAGCTTGA GCTTAAAGGA ACTTCTGATA AAAACAATGG ATCTGGAGTA  
420



-38-

CTTGAAGGCG TAAAAGCTGA CAAAAGTAAA GTAAAATTAA CAATTCTGA CGATCTAGGT  
480

CAAACACAC TTGAAGTTTT CAAAGAAGAT GGCAAAACAC TAGTATCAAA AAAAGTAACT  
540

TCCAAAGACA AGTCATCAAC AGAAGAAAAA TTCAATGAAA AAGGTGAAGT ATCTGAAAAA  
600

ATAATAACAA GAGCAGACGG AACCAGACTT GAATACACAG AAATTAAAAG CGATGGATCT  
660

GGAAAAGCTA AAGAGGTTTT AAAAAGCTAT GTTCTTGAAG GAACTTTAAC TGCTGAAAAA  
720

ACAACATTGG TGGTTAAGA AGGAACTGTT ACTTTAAGCA AAAATATTTT AAAATCTGGG  
780

GAAGTTTCAG TTGAACTTAA TGACACTGAC AGTAGTGCTG CTAATAAAAA AACTGCAGCT  
840

TGGAATTCAG GCACTTCAAC TTTAACAATT ACTGTAAACA GTAAAAAAC TAAAGACCTT  
900

GTGTTTACAA AAGAAACAC AATTACAGTA CAACAATACG ACTCAAATGG CACCAAATTA  
960

GAGGGGTCAG CAGTTGAAAT TACAAACTT GATGAAATTA AAAACGCTTT AAAATAA  
1017

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 338 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: BORRELIA BURGDORFERI  
 (B) STRAIN: ZS7

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: NS1-ZS7

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met	Asp	Pro	Asn	Thr	Val	Ser	Ser	Phe	Gln	Val	Asp	Ser	Phe	Leu	Trp
1				5					10					15	
His	Val	Arg	Lys	Arg	Val	Ala	Asp	Gln	Glu	Leu	Gly	Asp	Ala	Pro	Phe
			20					25					30		
Leu	Asp	Arg	Leu	Arg	Arg	Asp	Gln	Lys	Ser	Leu	Arg	Gly	Arg	Gly	Ser
			35					40					45		
Thr	Leu	Gly	Leu	Asp	Ile	Glu	Thr	Ala	Thr	Arg	Ala	Gly	Lys	Gln	Ile
			50					55					60		
Val	Glu	Arg	Ile	Leu	Lys	Glu	Glu	Ser	Asp	Glu	Ala	Leu	Lys	Met	Thr
			65					70					75		80
Met	Gly	Lys	Gln	Asn	Val	Ser	Ser	Leu	Asp	Glu	Lys	Asn	Ser	Val	Ser
				85						90				95	

-39-

Val Asp Leu Pro Gly Glu Met Asn Val Leu Val Ser Lys Glu Lys Asn  
 100 105 110  
 Lys Asp Gly Lys Tyr Asp Leu Ile Ala Thr Val Asp Lys Leu Glu Leu  
 115 120 125  
 Lys Gly Thr Ser Asp Lys Asn Asn Gly Ser Gly Val Leu Glu Gly Val  
 130 135 140  
 Lys Ala Asp Lys Ser Lys Val Lys Leu Thr Ile Ser Asp Asp Leu Gly  
 145 150 155 160  
 Gln Thr Thr Leu Glu Val Phe Lys Glu Asp Gly Lys Thr Leu Val Ser  
 165 170 175  
 Lys Lys Val Thr Ser Lys Asp Lys Ser Ser Thr Glu Glu Lys Phe Asn  
 180 185 190  
 Glu Lys Gly Glu Val Ser Glu Lys Ile Ile Thr Arg Ala Asp Gly Thr  
 195 200 205  
 Arg Leu Glu Tyr Thr Glu Ile Lys Ser Asp Gly Ser Gly Lys Ala Lys  
 210 215 220  
 Glu Val Leu Lys Ser Tyr Val Leu Glu Gly Thr Leu Thr Ala Glu Lys  
 225 230 235 240  
 Thr Thr Leu Val Val Lys Glu Gly Thr Val Thr Leu Ser Lys Asn Ile  
 245 250 255  
 Ser Lys Ser Gly Glu Val Ser Val Glu Leu Asn Asp Thr Asp Ser Ser  
 260 265 270  
 Ala Ala Thr Lys Lys Thr Ala Ala Trp Asn Ser Gly Thr Ser Thr Leu  
 275 280 285  
 Thr Ile Thr Val Asn Ser Lys Lys Thr Lys Asp Leu Val Phe Thr Lys  
 290 295 300  
 Glu Asn Thr Ile Thr Val Gln Gln Tyr Asp Ser Asn Gly Thr Lys Leu  
 305 310 315 320  
 Glu Gly Ser Ala Val Glu Ile Thr Lys Leu Asp Glu Ile Lys Asn Ala  
 325 330 335  
 Leu Lys

TABLE I

Strain	Origin	Source	PCR		Southern I Hind III * Osp A ( )
			aa 6-170 Osp A1-A4	aa 6-220 Osp A1-A2	
B31	USA	Ixodes dammini	+	+	I (A)
ZS7	Germany	I. ricinus	+	+	I (A)
Z37	Germany	I. ricinus	+	+	I (A)
GeHo	Germany	Skin (ECM)	+	+	I (A)
B1	Germany	Human Skin Biopsy	+	+	I (A)
B2	Germany	Human Skin Biopsy	+	+	I (A)
B3	Germany	Human Skin Biopsy	+	+	I (A)
20004	France	I. ricinus	+	+	I (A)
19535	USA	Peromyscus leucopus	+	+	I (A)
26816	USA	Microtus+	+	+	I (A)
28691	USA	I. dammini	+	+	I (A)
21305	USA	Peromyscus leucopus	+	+	I (A)
21343	USA	Peromyscus leucopus	+	+	I (A)
26815	USA	Chipmunk	+	+	I (A)
R7NE4	Switzerland	I. ricinus	+	+	I (A)
297	USA	Human/Spinal fluid	+	+	I (A)
MAC3	USA	Human	+	+	I (A)
20001	France	I. ricinus	+	+	I (A)
CT1P7	USA	Tick Isolate/dog	+	+	I (A)
SH-2-82-P5	USA	I. dammini	+	+	I (A)
CA-2-87	USA	I. pacificus	n.d.	n.d.	I (A)
S12/14	Germany	I. ricinus	+	+	I (A)
19857	USA	Rabbit Kidney	+	-	III (C)
ZQ1	Germany	I. ricinus	+	-	II (B)

TABLE I (Continued)

Strain	Origin	Source	PCR		Southern Hind III Osp A
			aa 6-170 Osp A1-A4	aa 6-220 Osp A1-A2	
NE4	Switzerland	I. ricinus	+	-	II (B)
NE58	Switzerland	I. ricinus	+	-	II (B)
NE11H	Switzerland	I. ricinus	+	n.d.	II (B)
R7NE58	Switzerland	I. ricinus	+	n.d.	II (B)
20047	France	I. ricinus	+	-	II (B)
N34	Germany	I. ricinus	+	-	II (B)
21038	USA	I. dentatus	+	-	III (C)
ACA-1	Sweden	Human skin	-	-	IV (D)
20047	France	I. ricinus	n.d.	n.d.	V (E)
S90	Germany	I. ricinus	n.d.	n.d.	VI (F)

\* alternative nomenclature

I Osp A genotype

- 42 -

Table II

## Crossprotection

immune serum transferred anti-	strain inoculated	protection (no arthritis)
ZS7	ZS7	+
ZS7	NE11H	-
NE11H	NEW11H	+
NE11H	ZS7	-
NE11H	20047 <sup>0</sup>	±
NE11H	19857 <sup>0</sup>	±
ZS7	20047 <sup>0</sup>	±
ZS7	19857 <sup>0</sup>	±
Cocktail*	20047 <sup>0</sup>	+
Cocktail*	19857 <sup>0</sup>	±

5

\* including 20047, ZS7 and NE11H;

0 strains showed reduced pathogenicity in unprotected controls.

- 43 -

**Table III - Crossprotection**

immune serum transferred anti-	strain inoculated	clinical arthritis d23-25p.i.	protection histo- pathology+	spirochetemia§
-	ZS7	8/8	+	5/7
ZS7	ZS7	0/3	-	0/3
ZS7	NE11H	3/3	+	3/3
ZS7	20047	(2/3) <sup>°</sup>	±	3/3
B31	ZS7	0/3	-	3/3
-	NE11H	7/7	+	4/5
NE11H	NE11H	(2/6) <sup>°</sup>	-^	0/6
NE11H	ZS7	3/3	+	3/3
NE11H	20047	(2/3) <sup>°</sup>	±	1/3
ZQ1	ZS7	3/3	+	3/3
-	20047	4/4	+	2/3
Cocktail*	20047	0/3	-^	1/3
rec.OspA	ZS7	(2/6) <sup>°</sup>	-^	5/6
rec.OspA	NE11H	3/3	+	2/3
rec.OspA	20047	(1/3) <sup>°</sup>	±	0/2
-	SH-2-82-P5	2/2	+	1/2
rec.OspA	SH-2-82-P5	(2/3) <sup>°</sup>	-^	0/3
-	21038	(2/2)	±	1/2 <sup>++</sup>
rec.OspA	21038	(2/3) <sup>°</sup>	-^	0/2

**Legend: Table III: Crossprotection**\* including *B.burgdorferi* organisms of strains 20047, ZS7 and NE11H;

5 + arthritis, carditis, hepatitis, myositis

° only, mild clinical arthritis

^ only minor histopathological alterations

§ recultivation

++ non-motil spirochaetes

Tabl IV

B. burgdorferi isolate	Biological origin	Geographical origin	fla/HSP60/HSP70 genogroup	OspA genotype
B31 (ATCC35210)	Tick (I. dammini)	USA	A/A/A	I
ZS7	Tick (I. ricinus)	Germany	A/A/A	I
Z37	Tick (I. ricinus)	Germany	A/A/A	I
GeHo	Skin (ECM)	Germany	A/A/A	I
B1	Skin (ECM)	Germany	A/A/A	I
B2	Skin (ECM)	Germany	A/A/A	I
B3	Skin (AD)	Germany	A/A/A	I
20004	Tick (I. ricinus)	France	A/A/A	I
19535	Mouse (Peromyscus)	USA	A/A/A	I
26816	Vole (Microtus)	USA	A/A/A	I
28691	Tick (I. dammini)	USA	A/A/A	I
21305	Mouse (Peromyscus)	USA	A/A/A	I
21343	Mouse (Peromyscus)	USA	A/A/A	I
26815	Chipmunk	USA	A/A/A	I
297	Cerebrospinal fluid	USA	A/A/A	I
Mac3	Skin	USA	A/A/A	I
20001	Tick (I. ricinus)	France	A/A/A	I
CTIP7	Dog tick	USA	A/A/A	I
SH-2-82	Tick (I. dammini)	USA	A/A/A	I
CA-2-87	Tick (I. pacificus)	USA	A/A/A	I
S12/14	Tick (I. ricinus)	USA	A/A/A	I
Z25	Tick (I. ricinus)	Germany	A/A/A	I
Z118	Tick (I. ricinus)	Germany	A/A/A	I

B. burgdorferi isolate	Biological origin	Geographical origin	fla/HSP60/HSP70 genogroup	OspA genotype
Z136	Tick (I. ricinus)	Germany	A/A/A	I
Z160	Tick (I. ricinus)	Germany	A/A/A	I
IP1	Cerebrospinal fluid	France	A/A/A	I
NE2	Tick (I. ricinus)	Switzerland	A/A/A	I
R7NE4	Tick (I. ricinus)	Switzerland	A/A/A	I
LW2	Skin	Germany	A/A/A	I
LW2.4	Skin	Germany	A/A/A	I
19857	Rabbit Kindey	USA	B/A/B/A	III
21038	Larva (I dentatus)	USA	B/A/B/A	III
ACA-1	Skin (ACA)	Sweden	B/B/A	IV
Bo23	Skin (ECM)	Germany	B/B/A	IV
So2	Tick (I. ricinus)	Great Britain	B/B/A	IV
PKo	Skin (ECM)	Germany	B/B/A	IV
ZQ1	Tick (I. ricinus)	Germany	B/B/B	II
NE4	Tick (I. ricinus)	Switzerland	B/B/B	II
NE58	Tick (I. ricinus)	Switzerland	B/B/B	II
NE11H	Tick (I. ricinus)	Switzerland	B/B/B	II
R3NE2	Tick (I. ricinus)	Switzerland	B/B/B	II
N34	Tick (I. ricinus)	Germany	B/B/B	II
IP3	Cerebrospinal fluid	France	B/B/B	II
IRS Us		Germany	B/B/B	II



B. burgdorferi isolate	Biological origin	Geographical origin	fla/HSP60/HSP70 genogroup	OspA genotype
Frst1	Skin	Germany	B/B/B	II
Frst2	Skin	Germany	B/B/B	II
So1	Tick (I. ricinus)	Great Britain	B/B/B	II
42/87		Sweden	B/B/B	II
152/86		Sweden	B/B/B	II
MK5	Tick (I. ricinus)	Hungary	B/B/B	II
MK6	Tick (I. ricinus)	Hungary	B/B/B	II
387	Cerebrospinal fluid	Germany	B/B/B	II
50	Cerebrospinal fluid	Germany	B/B/B	II
20047	Tick (I. ricinus)	France	B/B/B	V
S90	Tick (I. ricinus)	Germany	B/B/B	VI

**Claims**


1. A DNA sequence encoding an OspA protein or derivative thereof  
5 wherein the sequence encoding the OspA portion is substantially identical to the sequence depicted in Sequence ID No. 1 or 3.
2. A DNA sequence as claimed in claim 1 which is at 90% identical to the sequence depicted in Sequence ID No. 1 or 3.
- 10 3. A DNA sequence encoding an OspA derivative as claimed in claim 1 or 2 wherein the derivative is an NS1 - fusion protein.
4. A DNA sequence encoding an OspA protein characterised in that  
15 polymerase chain reaction will take place utilising prOspA1-A4 primers, but not prOsp A1-A2 primers.
5. An OspA protein or derivative thereof containing an amino acid  
20 sequence substantial identical to the sequence set forth in Sequence ID No. 2 or 4.
6. An OspA protein or derivative thereof which fails to react to monoclonal antibody LA2, but will react with monoclonal antibody LA31.
- 25 7. An OspA protein or derivative derived from B. burgdorferi subgroup II, III, IV V or VI.
8. An OspA derivative which is an NS1-fusion protein.
- 30 9. An NS1-OspA fusion protein wherein the OspA portion is selected from an OspA derived from B.burgdorferi species ZS7, ZQ1, 19857, and ACA I.
10. An NS1-OspA fusion protein as claimed in claim 8 or 9 wherein the  
35 OspA protein part is truncated.
11. A vaccine or diagnostic composition comprising at least two OspA proteins or derivatives thereof, characterised in that the OspA proteins or

- 48 -

derivatives are from different groups.

12. A vaccine composition comprising an OspA or derivative thereof, selected from the group, group I, group II, group III and group IV and a second OspA or derivative selected from the group I, group II, group III, and group IV provided that the first and second OspA are selected from a different group.
13. A vaccine as claimed in claim 11 or 12 comprising a ZS7 OspA or derivative thereof and a ZQ1 OspA or derivative thereof in admixture with a suitable carrier.
14. A vaccine composition comprising an OspA protein or derivative according to claim 5 or 6 in admixture with a suitable carrier.
15. A vaccine composition comprising an NS1-OspA fusion protein in admixture with a suitable carrier.
16. A vector comprising a DNA sequence as claimed in claim 1 to 4.
17. A method of producing an OspA protein or derivative thereof as claimed in claim 5 or 6, the method comprising:
- a) transforming a host with a vector of claim 16
  - b) expressing said protein or derivative in a suitable host
  - c) isolating the protein or derivative produced.
18. A method of treating a mammal susceptible of infection by *Borrelia* comprising administering an effective amount of a protein or derivative as claimed herein to the mammal.
19. A method of treating a mammal susceptible to infection by *Borrelia* comprising administering an effective amount of a vaccine as claimed herein.
20. A protein as claimed in any of claims 5 - 10 for use in medicine.
21. B. burgdorferi strain ZQ1.

1/1




SUBSTITUTE SHEET

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 92/01827

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/31; A61K39/02;	C12N15/44; C12N1/20;	C12N15/62; //C12P21/08 C07K13/00
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	C07K	
Documentation Searched other than Minimum Documentation to the extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
P, X	GENESEQ PROTEIN DATABASE, ACCESSION NO. X66065, DERWENT PUBLICATIONS LTD 3 June 1992, LONDON, GB R. WALLICH ET AL. 'Molecular heterogeneity among B. burgdorferi isolates' see the whole document	1, 4
Y	EP, A, 0 418 827 (MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V.) 27 March 1991 cited in the application	6, 7, 17, 18-20
A	see claims 1-3, 13, 15-25	3, 8-10, 15
	- / - -	
<p><sup>10</sup> Special categories of cited documents : <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
11 NOVEMBER 1992	24. 11. 92	
International Searching Authority	Signature of Authorized Officer	
EUR PEAN PATENT FFICE	THIELE U.H.-C.H. 	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y,0	<p>ABSTRACTS OF THE ANNUAL MEETING OF THE AMERICAN SOCIETY FOR MICROBIOLOGY, ABSTRACT NO. B-238 vol. 91, no. 0, 1991, WASHINGTON US page 65 S. BERGSTRÖM ET M. JONSSON 'Molecular characteristics and genetic organization of the osp-operon from different borrelia-burgdorferi isolates' see the whole document &amp; '91st General Meeting of the American Society for Microbiology 1991, Dallas, Texas, USA, May 5-9, 1991'</p> <p>---</p>	6,7,17, 18-20
X	<p>DER HAUTARZT vol. 41, no. 12, 1990, BERLIN, FRG pages 648 - 657 M. D. KRAMER ET AL. 'Die Borrelia-burgdorferi-Infektion'</p> <p>---</p>	21
A	<p>see page 654, left column, paragraph 3</p> <p>---</p>	6,7,20
A	<p>EP,A,0 366 238 (SMITHKLINE BEECHAM CORPORATION) 2 May 1990 see claims 1,3,4</p> <p>-----</p>	3,8-10, 15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 92/01827

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 18,19 are directed to a method of treatment of the human/animal body (see PCT-Rule 39.1(iv)) the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO. EP 9201827  
SA 63773**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 11/11/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0418827	27-03-91	DE-A- 4015911	28-03-91
		AU-A- 6244490	28-03-91
		CA-A- 2025597	20-03-91
		JP-A- 3209400	12-09-91
EP-A-0366238	02-05-90	AU-A- 4022489	15-03-90
		JP-A- 3130082	03-06-91

EPO FORM P0019

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82